

# Coadministración de vacunas en preadolescentes

30/01/2018

**Respuesta del Experto a ...**

Coadministración de vacunas en preadolescentes

## **Pregunta**

Con la introducción de calendario vacunal 2018 en Andalucía , se aconseja que a la hora de administrar VARICELA tanto con VPH como con Meningococo C , a jóvenes de 12-14 años se espacie la administración 1 mes entre ambas , pero no encuentro evidencia de ello. ¿ Existe alguna evidencia del dicho consejo, puede existir interferencia inmunógena ?

## **Respuesta de José Antonio Navarro (30 de Enero de 2018)**

Los estudios de inmunogenicidad en niños y/o preadolescentes respecto a la administración concomitante de la vacuna VPH junto a varicela y antimeningocócica conjugada tetravalente (CRM) no han mostrado interferencias en la respuesta inmune a todas ellas <sup>(1, 2)</sup>.

## **Referencias**

<sup>(1)</sup> Gasparini R et al. Safety and Immunogenicity of a Quadrivalent Meningococcal Conjugate Vaccine and Commonly Administered Vaccines After Coadministration. *Pediatr Infect Dis J* 2016;35:81-93

<sup>(2)</sup> Abdelnour A et al. Safety of a quadrivalent meningococcal serogroups A, C, W, and Y conjugate vaccine (MenACWY-CRM) administered with routine infant vaccinations: Results of an open-label, randomized, phase 3b controlled study in healthy infants. *Vaccine* 2014;32:965-972